

# Sound-induced monoaminergic turnover in the auditory forebrain depends on endocrine state in a seasonally-breeding songbird

C. A. Rodríguez-Saltos<sup>1</sup>  | S. M. Lyons<sup>2</sup> | K. W. Sockman<sup>2</sup> | D. L. Maney<sup>1</sup> 

<sup>1</sup>Department of Psychology, Emory University, Atlanta, GA, USA

<sup>2</sup>Department of Biology, University of North Carolina, Chapel Hill, NC, USA

## Correspondence

Donna L. Maney, Department of Biology, Rollins Research Center, Emory University, Atlanta, GA, USA.  
Email: dmaney@emory.edu

## Funding information

Howard Hughes Medical Institute; National Science Foundation, Grant/Award Number: NSF-IOS 0723805; National Institutes of Health, Grant/Award Number: 5R01NS055125-04

Sensory responses to courtship signals can be altered by reproductive hormones. In seasonally-breeding female songbirds, for example, sound-induced immediate early gene expression in the auditory pathway is selective for male song over behaviourally irrelevant sounds only when plasma oestradiol reaches breeding-like levels. This selectivity has been hypothesised to be mediated by the release of monoaminergic neuromodulators in the auditory pathway. We previously showed that in oestrogen-primed female white-throated sparrows, exposure to male song induced dopamine and serotonin release in auditory regions. To mediate hormone-dependent selectivity, this release must be (i) selective for song and (ii) modulated by endocrine state. Therefore, in the present study, we addressed both questions by conducting playbacks of song or a control sound to females in a breeding-like or a nonbreeding endocrine state. We then used high-performance liquid chromatography to measure turnover of dopamine, norepinephrine and serotonin in the auditory midbrain and forebrain. We found that sound-induced turnover of dopamine and serotonin depended on endocrine state; hearing sound increased turnover in the auditory forebrain only in the birds in a breeding-like endocrine state. Contrary to our expectations, these increases occurred in response to either song or artificial tones; in other words, they were not selective for song. The selectivity of sound-induced monoamine release was thus strikingly different from that of immediate early gene responses described in previous studies. We did, however, find that constitutive monoamine release was altered by endocrine state; irrespective of whether the birds heard sound or not, turnover of serotonin in the auditory forebrain was higher in a breeding-like state than in a nonbreeding endocrine state. The results of the present study suggest that dopaminergic and serotonergic responses to song and other sounds, as well as serotonergic tone in auditory areas, could be seasonally modulated.

## KEYWORDS

catecholamine, dopamine, norepinephrine, serotonin, song, white-throated sparrow


## 1 | INTRODUCTION

Courtship signals command the attention of their intended receivers.<sup>1-4</sup> Because courtship signals are relevant only to receivers in breeding condition, they should command more attention during the breeding season than outside it. In seasonally-breeding female *Zonotrichia* sparrows, breeding-typical plasma levels of oestradiol are required for proceptive responses to playback of male song.<sup>5-7</sup> Because behavioural responses to courtship signals depend on endocrine state, we expect that the neural mechanisms underlying the salience of those signals depend on endocrine state as well. In a variety of seasonally breeding songbirds, auditory responses to conspecific song are in fact greater during the breeding season. Such changes have been demonstrated using auditory evoked potentials,<sup>8,9</sup> neurophysiological recording,<sup>10,11</sup> in vivo imaging<sup>12-14</sup> and the expression of immediate early genes.<sup>5,15</sup> There is thus good evidence that the auditory system becomes more attuned to conspecific song when a bird enters a breeding-like endocrine state.

The brain ascribes salience to sensory signals in part via the monoaminergic neuromodulators dopamine, norepinephrine and serotonin.<sup>2</sup> These systems are evolutionarily ancient and can be found in all vertebrate taxa.<sup>16</sup> Fibres carrying these neuromodulators originate in a small number of nuclei but project widely throughout the brain. Each of the monoaminergic systems serves multiple functions. In the present study, we focus on their shared function of modulating auditory responses. Noradrenergic, dopaminergic and serotonergic fibres terminate at all levels of the auditory system in vertebrates<sup>17</sup> and facilitate attention to and learning of auditory stimuli.<sup>2</sup>

In birds, monoaminergic cell bodies project from the brainstem to almost every area of the brain, including auditory areas.<sup>18-20</sup> Lesions of brainstem noradrenergic neurones inhibited behavioural responses to song in female canaries<sup>21, cf 22</sup> and eliminated behavioural preferences for more attractive vs less attractive song in female zebra finches.<sup>23</sup> Noradrenergic receptor antagonism in European starlings disrupted female preferences for conspecific over heterospecific song<sup>24</sup> and treatment with a dopamine agonist disrupted preferences for attractive song.<sup>25</sup> In zebra finches, playback of novel songs induced more immediate early gene expression in noradrenergic cell bodies than playback of familiar song, and this induction was correlated with induction in the auditory forebrain, particularly in cells contacted by catecholaminergic fibres.<sup>26</sup> Taken together, these studies suggest that hearing song engages monoaminergic systems, which may be necessary for appropriate auditory and behavioural responses.

Because monoaminergic systems are highly sensitive to reproductive hormones, they are prime candidates for seasonal modulation of behavioural responses.<sup>2,27</sup> The transition to breeding, accompanied by increases in plasma sex steroids, may bring increases in constitutive monoamine release that persist over weeks, priming the auditory pathway to respond to courtship signals.<sup>2</sup> In female white-throated sparrows, treatment with oestradiol increased the density of both catecholaminergic and serotonergic fibres at multiple levels of the auditory pathway.<sup>19,20,28</sup> Oestradiol could therefore effectively increase the amount of catecholamine or serotonin available for release in response to song. Alternatively, or in addition, oestradiol may increase the rate of tonic or nonsynaptic release that occurs independently of sensory stimulation. In rats,



**FIGURE 1** Experimental design. The aims of the present study were 2-fold (A). We previously showed that in oestradiol-primed female white-throated sparrows, hearing male song causes release of dopamine and serotonin in the auditory forebrain.<sup>20,30</sup> Question 1 in the present study is whether that release depends on endocrine state and is selective for song. Question 2 is whether monoamine turnover increases in a breeding-like endocrine state, independent of hearing sound, which may prime the auditory system to respond selectively to song. We tested these hypotheses using the design depicted in (B). Sixty animals were collected and housed in aviaries on short days before being transferred to individual cages and housed 5 cages per small room. Each of the rooms contained 1 bird that would hear one of the 5 types of sound stimulus. Half of the birds remained on a winter-like photoperiod (blue) and half were photostimulated (red). After 3 weeks of individual housing, birds received s.c. silastic capsules containing nothing (blue) or oestradiol (red). One week later, each bird was isolated in a sound-attenuated booth and, the next morning, it was presented with a sound stimulus. Brains were collected and frozen immediately following the stimulus presentation. Brains were sectioned sagittally at 300  $\mu$ m and regions of interest microdissected as shown by the black circles in (C). CMM, caudomedial mesopallium. IC, inferior colliculus. NCM, caudomedial nidopallium. For further details, see Materials and Methods

stimulus-induced and stimulus-independent release of dopamine occur even in the same population of neurones.<sup>29</sup> These actions of monoamines (ie, induced by stimuli or via priming) (Figure 1A) are thus not mutually exclusive.

In a previous study of female white-throated sparrows, playback of male conspecific song induced release of both dopamine and serotonin in the auditory forebrain.<sup>20,30</sup> This release was detectable, using high-performance liquid chromatography (HPLC) of micropunched tissue, after 30 minutes of song exposure. Thus, there is evidence that hearing song can induce monoamine release fairly quickly in an auditory area. Those studies left several questions unanswered. Because all of the birds in those studies were treated with oestradiol, it is not known whether the song-induced monoamine release depended on endocrine state. In addition, because no control sounds were presented, it is not known whether the monoamine release was caused by exposure to song specifically, or whether it would have occurred in response to any sound. In the present study, we manipulated endocrine state in female white-throated sparrows and then exposed them to conspecific male song, a control sound or silence. Following stimulus presentation, we measured turnover of dopamine, norepinephrine and serotonin in the auditory midbrain and 2 regions of the auditory forebrain. We predicted that if neural responses to song are driven or modulated by monoamine release on a rapid timescale, then monoamine turnover in these areas would be higher in birds hearing song than in birds hearing tones or silence. In addition, we predicted that such effects would be apparent only in birds in a breeding-like (oestradiol-primed and photostimulated) endocrine state. Finally, we predicted that if the constitutive release of monoamines depends on endocrine state, turnover would be higher in birds in breeding-like condition than in birds in nonbreeding condition, regardless of the type of stimulus presented. These results would support our hypothesis that the salience of male song, which attracts females only during the breeding season, is mediated by hormone-dependent monoamine release within the auditory system.

## 2 | MATERIALS AND METHODS

### 2.1 | Animals

Animals were collected under appropriate local, state and federal permits. Emory University's Institutional Animal Care and Use Committee approved all of the procedures that involved the use of animals. Sixty female white-throated sparrows (*Zonotrichia albicollis*) were trapped using mist-nets around Emory University campus during their autumn migration. After capture, the birds were housed in indoor aviaries in the Division of Animal Resources at Emory University. The birds were held under a winter-like photoperiod (9.5:14.5 hour light/dark photoperiod) until at least mid-January so that they would regain photosensitivity.<sup>31,32</sup> Housing under a short photoperiod prevents increases in endogenous plasma oestradiol in this species.<sup>5,6</sup> Food and water were provided ad libitum.

The sex of the birds was determined by polymerase chain reaction (PCR) analysis<sup>33</sup> of a blood sample obtained by venipuncture of the brachial vein. Sex was then confirmed at the end of the study by inspection of the gonads during necropsy. Age and plumage morph, which could affect behavioural responses to song playback,<sup>34</sup> were balanced across our experimental treatments (age:  $n = 23$  hatch-years,  $n = 8$  after hatch-years,  $n = 5$  unknown age; morph:  $n = 15$  tan-striped,  $n = 21$  white-striped). Age was determined by non-invasive inspection of plumage and skulls in live birds.<sup>35</sup> Morph was determined by PCR analysis.<sup>36</sup>

### 2.2 | Manipulation of endocrine state

Our experimental design is depicted in Figure 1(B). All birds were transferred to individual cages and housed in small rooms ( $n = 5$  per room). We assigned the birds to 1 of 2 treatment groups. For birds in the first group ( $n = 30$ ), the light cycle was changed to a summer-like photoperiod (14:10 hour light/dark photoperiod) to facilitate a breeding-like endocrine state. Because photostimulation alone is not sufficient to elevate plasma oestradiol to levels typical of the reproductive season,<sup>6,7,37</sup> after 3 weeks of photostimulation, we treated birds in this group with exogenous oestradiol via a s.c. implant.<sup>19</sup> Each implant consisted of a silastic capsule (length 12 mm, inner diameter 1.47 mm, outer diameter 1.96 mm; Dow Corning, Midland, MI, USA) filled with  $17\beta$ -oestradiol (Steraloids, Newport, RI, USA) and sealed at both ends with silicon adhesive. This dose has been demonstrated to increase plasma oestradiol to breeding-typical levels in captive females of this genus after 2 days of treatment for up to 80 days.<sup>7</sup> In this species, this dose brings plasma oestradiol levels up to  $1.3 \pm 0.2$  ng mL<sup>-1</sup>,<sup>5</sup> which is within the physiological range for free-living, breeding females early in the breeding season when they are likely attending to male song.<sup>38</sup> We have previously shown, in multiple studies, that this dose of oestradiol increases the selectivity of immediate early gene responses in the auditory forebrain and midbrain.<sup>5,6,39,40</sup> Moreover, hearing song increased concentrations of dopamine and serotonin metabolites in the auditory forebrain in females treated with this dose.<sup>20,30</sup>

The birds in the second group ( $n = 30$ ) were kept in a non-breeding endocrine state. These birds continued to be held on a winter-like photoperiod (9.5:14.5 hour light/dark cycle) and received empty silastic capsules. Under these conditions, ovaries remain regressed and plasma oestradiol remains very low.<sup>5,6</sup> After completion of the study, ovaries were inspected to confirm a regressed state.

### 2.3 | Playback experiment

Four weeks following the onset of the above treatment (and 1 week following implant administration), we exposed each bird to 1 of 5 auditory stimuli: silence, conspecific male song for 15 or 30 minutes or artificial tones for 15 or 30 minutes. Six birds in each of the 2 endocrine states heard each type of stimulus. Type of stimulus was

balanced across birds that were housed in the same room during the manipulation of breeding state, so that each of the 5 birds in a room was exposed to a different stimulus during the playback experiment (Figure 1B). The sound presentations have been described elsewhere.<sup>5,20,30,41</sup> Briefly, songs from several male white-throated sparrows were downloaded from the Borror Bioacoustic Library (<https://blb.osu.edu/>). Using AUDACITY (<https://www.audacityteam.org>), we constructed sound presentations in which a song from a unique male was repeated every 15 seconds and the identity of the singer changed every 3 minutes. Thus, each 15-min presentation contained 5 different male songs, each repeated 12 times, and each 30-min presentation contained 10 male songs each repeated 12 times. Six versions of each presentation were constructed, 1 for every bird in each treatment group assigned to hear song. The versions differed in the order in which the songs were presented, following a Latin-square design.

For each song, a sequence of artificial tones was generated to match the dominant frequencies in the song.<sup>5</sup> The order of the tones within each sequence was random and their durations equal. The overall duration of the tone sequence equaled that of the song. The tone sequences were automatically generated with a script (available upon request) written in R (R Core Team, 2013), using functions from the software packages *seewave*<sup>42</sup> and *tuneR*.<sup>43</sup> The tone sequences were arranged into stimulus presentations exactly as described above for the songs such that, for each of the 6 unique song presentations, there was a matching tone presentation. As was the case for the birds hearing song, each of the 6 birds per experimental group heard a unique tone presentation.

On the afternoon before the playback, each bird was isolated in a sound-attenuating booth (Industrial Acoustics, Bronx, NY, USA). The booth was equipped with a microphone, a speaker (LS-300; AudioSource, Portland, OR, USA) and a video camera (WV-CP240; Panasonic, Osaka, Japan) that recorded video into a digital video recorder (DVR 1604 HF-U; Dahua, Hangzhou, China). The bird remained isolated in the booth overnight, and presentation of conspecific song, tones or silence (no sound stimulus) started between 30 and 75 minutes after lights on (between 08.00 and 08.45 hours). Bird behaviour was recorded on video throughout the stimulus presentation. Birds assigned to the silence group remained in the booth for the same amount of time as did the birds that heard song or tones.

## 2.4 | Tissue collection

For birds that heard song or tones, brains were collected immediately after playback. For birds that heard silence, brains were collected within 60–100 minutes after lights on such that they were collected at the same time of day as were the other brains. Brains were rapidly frozen in powdered dry ice within 10 minutes after rapid decapitation. Sagittal frozen sections (300  $\mu\text{m}$ ) were mounted on microscope slides by rapid thawing and refreezing. We used the “Palkovits Punch Technique” [44] to extract brain tissue from 3

auditory regions of interest (ROIs) as depicted in Figure 1(C): the caudomedial mesopallium (CMM) and caudomedial nidopallium (NCM), which are regions of the auditory forebrain homologous to secondary auditory regions of mammals,<sup>45,46</sup> and the inferior colliculus (IC; in birds also called the dorsal lateral nucleus of the mesencephalon). The diameter of the punches was 0.5 mm for CMM and IC, and 1.0 mm for NCM (Figure 1C). Two punches were collected for each ROI. We used tissue from either the right or the left hemisphere, balanced across groups.

## 2.5 | Measurement of monoaminergic activity

We measured the concentrations of the monoamines dopamine, norepinephrine and serotonin, and their respective metabolites dihydroxyphenylacetic acid (DOPAC), 3-methoxy-4-hydroxyphenyl ethylene glycol and 5-hydroxyindolacetic acid using HPLC with electrochemical detection.<sup>47</sup> Preparation of the samples and the mobile phase for HPLC followed protocols reported elsewhere.<sup>48</sup> HPLC was carried out using an HTEC 500 HPLC-ECD detector system (Eicom, Dublin, Ireland). Analytes were automatically identified using the software PowerChrome (edaa, Denistone East, NSW, Australia). The software matched the retention times of peaks in the HPLC output trace to the retention times of peaks corresponding to 1  $\mu\text{g } \mu\text{L}^{-1}$  and 10  $\mu\text{g } \mu\text{L}^{-1}$  external standards. A 5% error margin was allowed for matching retention times. When the software failed to recognise a peak, we classified that peak manually. The peak heights of the samples were normalised to that of an internal 10  $\mu\text{g } \mu\text{L}^{-1}$  isoproterenol standard. The concentration of the analytes was estimated from a linear interpolation between the concentrations and normalised peak heights of the external standards.

Because we were interested in quantifying monoaminergic activity rather than absolute concentration, we estimated turnover. Turnover is assumed to be related to the rate of release<sup>47</sup> and has been used to estimate such in songbirds.<sup>49–51</sup> To be consistent with this literature, we defined turnover of each monoamine as the ratio of each metabolite to the corresponding monoamine. In other words, dopamine turnover was calculated by dividing the DOPAC value by the dopamine value, and so on.

## 2.6 | Statistical analysis

Our primary goal was to replicate and extend the findings of Matragrano et al.<sup>20,30</sup>. Accordingly, we used a similar statistical approach. We performed a separate analysis for each of the 3 monoamine turnover rates (dopamine, norepinephrine and serotonin) in each of the 3 auditory regions (NCM, CMM and IC). Because Matragrano et al.<sup>20,30</sup> found effects of playback at 30 minutes, we were primarily interested in that time point and analysed those data first. To determine whether the response to hearing song depends on endocrine state (Figure 1A), we tested whether the difference between silence and song depended on treatment (Figure 1A, Question 1); in other words, we were interested in the interactions with endocrine state. To test for these

interactions, we used mixed effects linear models in which monoamine turnover was the response. Each of these models included the following predictors: treatment (breeding-like or nonbreeding); the contrast between silence and song (to test whether hearing song induced turnover); the contrast between song and tones (to test for selectivity of the response); and each 2-way interaction (to test whether either effect depended on endocrine state). To account for the non-independence of each group of individuals that were housed together in the same room (1 hearing silence, 1 hearing song for 30 minutes, and 1 hearing tones for 30 minutes; see Figure 1B), observation was nested within group as a random intercept and as a random coefficient for all predictors except treatment. This random effects structure followed that described by Schielzeth and Forstmeier.<sup>52</sup>

To gain insight about how rapidly effects could be seen, we then analysed the data from the 15-minute time point as described above; for that time point, the data from the birds hearing silence were re-used. As described above, non-independence of the individuals housed together (one hearing silence, 1 hearing 15 minutes of song, 1 hearing 15 minutes of tones) (Figure 1B) was accounted for.

When the contrast between song and silence interacted with treatment (ie, when the effect of hearing song depended on endocrine state), we followed up with post-hoc comparisons to test whether the response to song differed from the response to tones within each endocrine state, and whether endocrine state affected responses to each type of stimulus.

The statistical approach described above allowed us to ask both parts of Question 1 (Figure 1A). We next aimed to ask Question 2: whether endocrine state had a main effect on constitutive (stimulus-independent) monoamine turnover. In this test, we wanted to include all 60 of our experimental animals in the same model to maximise our power to detect a main effect of treatment. At the same time, we needed to control for the effects of hearing sound on monoamine turnover. To accomplish these goals, we fitted an additional set of mixed effects linear models, this time including all 60 birds in each model. Treatment was included as a predictor. In addition, to control for sound playback, we also included the independent contrasts between silence and each duration of song or tones. Interactions were not included in this model. For these analyses, we assessed only the effects of treatment.

To rule out copulation solicitation displays as a source of variation in monoaminergic activity, we tested for a linear correlation between the number of these displays and monoamine turnover. Displays were scored from the videos recorded during playback. Only data from oestradiol-treated birds were used because nonbreeding birds of this species are not known to perform courtship displays and none of the birds in nonbreeding condition did so in the present study. We included data from the birds in breeding-like condition that were exposed to song, tones or silence because the birds in our laboratory occasionally solicit spontaneously, regardless of any sound stimulus.

## 3 | RESULTS

### 3.1 | Sound-induced monoamine turnover


The effects of hearing sound on monoamine turnover are shown in Figure 2. The means and SEs are provided in the Supporting information (Table S1). The statistical models, including  $z$  and  $P$  values for all contrasts and post-hoc tests, are also provided in the Supporting information (Table S2).

To test the hypothesis that song-induced monoamine release depends on endocrine state, we were interested in the interaction between the effects of song (compared with silence) and endocrine state. We found such interactions after 30 minutes of song playback for dopamine turnover in NCM ( $z = 2.03$ ;  $P = .043$ ), as well as for serotonin turnover in NCM ( $z = 2.24$ ;  $P = .025$ ) and CMM ( $z = 2.30$ ;  $P = .021$ ). In none of these cases, however, did endocrine state interact with the contrast between song and tones (all  $P > .2$ ). In other words, we found no evidence that the selectivity of these responses depended on endocrine state. We found no interactions between the silence-song contrast and treatment, nor any main effects of song, in IC.

Post-hoc tests showed that, for birds in breeding-like condition, although hearing song induced turnover of both dopamine and serotonin, the turnover was not different from that after exposure to tones (Figure 2). For example, dopamine turnover in NCM was higher in birds hearing 30 minutes of song than in birds hearing silence ( $z = 4.88$ ,  $P < .001$ ) but not higher than in birds hearing 30 minutes of tones ( $z = 0.71$ ,  $P = .48$ ). Dopamine turnover in NCM was higher in the breeding-like birds than in the nonbreeding birds after presentation of either 30 minutes of song ( $z = 1.65$ ,  $P = .099$ , note trend) or 30 minutes of tones ( $z = 5.38$ ,  $P < .001$ ). The effect of hearing song on dopamine turnover in NCM was significant after hearing only 15 minutes of song ( $z = 2.52$ ,  $P = .012$ ) but, again, this response was not different from the response to tones ( $z = 0.32$ ,  $P = .747$ ). Overall, for birds in a breeding-like endocrine state, hearing sound did elicit dopamine release but it did not matter whether song or tones was played.

The results of post-hoc tests on serotonin turnover in NCM were similar to those for dopamine. Compared with silence, exposure to 30 minutes of song increased serotonin turnover in the birds in breeding-like condition ( $z = 2.58$ ,  $P = .01$ ). Song-induced turnover, however, was not higher than after exposure to 30 minutes of tones ( $z = 0.58$ ,  $P = .563$ ). As was the case for dopamine, serotonin turnover was higher in the breeding-like than in the nonbreeding birds after exposure to either song ( $z = 2.92$ ,  $P = .004$ ) or tones ( $z = 2.76$ ,  $P = .006$ ). Unlike song-induced dopamine turnover, serotonin turnover did not change by 15 minutes after onset of song playback ( $z = 0.74$ ,  $P = .457$ ).

Post-hoc tests on serotonin turnover in CMM revealed results identical to those in NCM. Within the birds in breeding-like condition, exposure to song for 30 minutes induced serotonin turnover in CMM ( $z = 2.20$ ,  $P = .028$ ). This response to song was not, however, different from the response to tones ( $z = 0.94$ ,  $P = .349$ ). As was the



**FIGURE 2** Effects of sound playback and endocrine state on monoamine turnover in the auditory forebrain. The effects of song exposure on dopamine and serotonin turnover depended on endocrine state. In birds that were photostimulated and treated with oestradiol (“breeding-like”; red lines), hearing song (solid lines) increased turnover of dopamine in NCM (A), serotonin in NCM (C) and serotonin in CMM (F) compared with silence (open triangles) ( $*P < .05$ ). These effects were, however, not different from the effect of exposure to tones (dotted lines). In birds in nonbreeding condition (blue lines), monoamine turnover was unaffected by exposure to either sound. The line graph in (A) excludes an extreme outlier that did not change the outcome of the statistical tests. Insets depict the main effects of endocrine state on turnover ( $*P < .05$ ), with all animals included in the same analysis and the effects of sound stimulus controlled. Estimated marginal means (ie, the means adjusted for the effects of sound stimulus) are shown for each endocrine state (blue, nonbreeding; red, breeding-like). IC, inferior colliculus; CMM, caudomedial mesopallium; NCM, caudomedial nidopallium

case for NCM, serotonin turnover in CMM was higher in the breeding-like than in the nonbreeding birds after exposure to either song ( $z = 3.11$ ,  $P = .002$ ) or tones ( $z = 2.72$ ,  $P = .007$ ). Serotonin turnover did not change after only 15 minutes of song playback ( $z = 0.74$ ,  $P = .457$ ). For both NCM and CMM, although exposure to 30 minutes of song increased serotonin turnover only in the birds in a breeding-like endocrine state, we found no evidence that this turnover was stimulated by song specifically instead of by sound in general.

### 3.2 | Effects of endocrine state on constitutive monoamine turnover

The main effects of endocrine state are shown in the insets in Figure 2. We found significant main effects of endocrine state on serotonin turnover in NCM ( $z = 3.61$ ;  $P < .001$ ) and CMM ( $z = 3.28$ ;  $P < .001$ ). No other significant main effects of endocrine state were detected. Overall, these results are consistent with the view that serotonergic tone in the auditory system is modulated seasonally in this species.

### 3.3 | Correlations with behaviour

To test whether proceptive behaviour could have contributed to increases in monoaminergic activity, we tested for correlations between the number of solicitation displays and monoaminergic activity in birds in the oestradiol-treated condition. We found no evidence of such a correlation. Spearman rho estimates were low overall, at between  $-0.240$  and  $0.194$  for all tests. Our ability to detect such correlations was limited by sample size because only the oestradiol-treated birds were expected to solicit<sup>7</sup> and, of those, only those exposed to song (as opposed to tones or silence) were expected to perform appreciable numbers of displays. In the present study, 2 birds exhibited 95% of all the displays that were recorded. Because a relatively low number of birds showed proceptive behaviour, it is unlikely that such behaviour contributed to the significant effects of sound or endocrine state on monoaminergic activity.

### 3.4 | Confirmation of ovarian regression

None of the birds in the unphotostimulated group showed any sign of ovarian development (very small ovary, largest ovarian follicle <1 mm). The birds in the photostimulated group showed signs of mild ovarian development; their ovaries and largest ovarian follicle were, on average, larger than those in the unphotostimulated group (Fisher's exact test,  $P < .0001$ ).

## 4 | DISCUSSION

In previous studies, we showed that hearing 30 minutes of male song caused an increase in dopamine and serotonin metabolites in NCM in oestrogen-primed female white-throated sparrows.<sup>20,30</sup> In the present study, we tested whether sound-induced increases in monoaminergic activity are higher in response to song than to a control sound, and whether they depend on endocrine state (Figure 1A, Question 1). On the basis of popular models, in which monoamines modulate auditory responses in accordance with the behavioural relevance of the stimulus,<sup>2,27,53-56</sup> we predicted that hearing song, but not a behaviourally irrelevant tone sequence, would increase monoaminergic turnover in the auditory pathway. Our data suggest that although dopamine and serotonin were released in NCM in response to hearing song (Figure 2), this release did not differ from the response to tones. Thus, the increases in dopaminergic and serotonergic activity described by Matrigrano et al.<sup>20,30</sup> may not have been driven specifically by song; the results of the present study suggest that they would have occurred in response to other sounds.

This result is surprising because songs are presumably more behaviourally relevant than our tone sequences. It is possible that for some reason, this control stimulus was perceived as particularly behaviourally relevant. Novel sounds induce greater immediate early gene expression in NCM than do familiar sounds,<sup>57-59</sup> and our birds were naïve to the tone sequences. We consider this explanation to be unlikely. We have previously shown, in multiple experiments in the same species, that these tones drive significantly lower Egr-1 (ZENK) responses in NCM, CMM, the auditory thalamus and IC than do conspecific songs.<sup>5,39,40,60</sup> The Egr-1 response in the auditory forebrain is so closely related to the behavioural relevance of the stimulus that it has been used as a proxy for such.<sup>57,61</sup> Egr-1 induction is also closely tied to neuromodulatory activity.<sup>2</sup> In zebra finches, application of norepinephrine directly to NCM induced Egr-1 expression<sup>62</sup> and, in European starlings, depletion of norepinephrine significantly attenuated the Egr-1 response to song.<sup>63,cf 22</sup> We therefore expected that sound-induced monoamine activity would mirror the previously described Egr-1 response, which is known to be selective for song over tones in all 3 auditory areas in the present study.<sup>39</sup> This prediction, however, was not borne out; the monoamine responses that we detected were not any higher to song than they were to tones. Thus, differential monoamine release is perhaps not a good candidate

mechanism for explaining the selectivity of Egr-1 responses to song.

Other playback studies have provided little evidence that monoaminergic neurones in the brainstem respond selectively to song. In the same birds that showed selective Egr-1 responses in auditory areas, we could not detect selective Egr-1 responses in dopaminergic, noradrenergic or serotonergic cell bodies.<sup>28,54</sup> Lynch et al.<sup>64</sup> found that in female canaries, hearing song induced Egr-1 expression in noradrenergic cells, but no control sounds were presented in that study, however. Chen et al.<sup>65</sup> found that in juvenile male zebra finches, Fos was induced in noradrenergic and dopaminergic cells, although only when song was presented by a live tutor, not by passive playback.<sup>cf 66, 67</sup> In all of the above studies, monoaminergic cell activation was determined by labelling immediate early gene expression, which may not reflect the relevant activity. Immediate early genes are not always induced by neuronal firing<sup>68</sup> and can usually be sampled at only 1 time point following the presentation of a stimulus in each animal. Using electrophysiological recording, Gale and Perkel<sup>69</sup> found that dopaminergic cells of the ventral tegmental area in zebra finches fired selectively in response to the bird's own song. Thus, it may be possible to detect song-selective activity of monoaminergic cell groups using more time-sensitive techniques.

Although monoamine release in the present study did not depend on behavioural relevance of the sound playbacks, it did depend on endocrine state. In NCM, sound-induced release of dopamine and serotonin was detected only in birds that had been photostimulated and treated with oestradiol. Thus, we replicated our previous finding of song-induced dopamine and serotonin release in oestradiol-treated birds,<sup>20,30</sup> and showed further that it does not occur in nonbreeding birds. Similarly, artificial tones induced release only in birds in breeding condition, suggesting that the reproductive endocrine state facilitates changes in the neurochemistry of the auditory forebrain that are relevant not just to conspecific vocalisations, but generally.

Increased monoamine release during the breeding season may be mediated by an effect of reproductive hormones on releasable stores.<sup>2</sup> In previous studies of female white-throated sparrows, oestradiol treatment increased the density of catecholaminergic and serotonergic innervation of CMM and NCM.<sup>19,20</sup> It is therefore possible that the enhanced sound-induced release that we detected in the birds in breeding-like condition can be explained simply by an increase in the availability of monoamines to be released. On a functional level, the result is harder to explain because it is not clear why both song and tone sequences would be more behaviourally relevant in spring than autumn. A more complete understanding of the monoaminergic responses will require investigating release in regions outside the auditory system. Behavioural responses to relevant sounds are not determined or orchestrated within any single system, particularly not the auditory system alone.<sup>27,54</sup> Other systems (eg, the reward pathway and hypothalamus) are involved and must also be examined.<sup>22,56,70,71</sup>

The results of the present study suggest that auditory forebrain selectivity for behaviourally relevant sounds cannot be

explained by event-related monoamine release during sound exposure. Monoamines could tune responses, however, by acting on a longer timescale. Monoaminergic neuromodulators are released from fibres via both synaptic and nonsynaptic mechanisms.<sup>72</sup> Norepinephrine applied directly to NCM in zebra finches increased the signal-to-noise ratio and the coding accuracy of single neurones,<sup>73</sup> suggesting permissive effects on the detection and encoding of complex sounds such as song. In the present study, photostimulation plus oestradiol administration did not increase noradrenergic turnover. Serotonergic tone, on the other hand, was dramatically increased in NCM and CMM. Thus, the answer to our Question 2 (Figure 1A) was that serotonergic activity in the auditory pathway does depend on endocrine state. These results are consistent with our previous findings that oestradiol administration increased the concentration of a serotonin metabolite in the auditory forebrain in white-throated sparrows.<sup>20</sup> These increases could serve as a mechanism by which oestradiol, working indirectly via serotonin, primes the auditory system to respond selectively to song.

Two potential caveats should be considered when interpreting our results. First, we measured monoamine turnover in dissected tissue, which does not afford the best time resolution. Methods of measuring release in real time, such as voltammetry,<sup>74-76</sup> should be employed to better detect event-related release. Second, our definition of monoamine turnover is based on the assumption that an increase in metabolites, relative to the concentration of the monoamines themselves, indicates release. Our results could also indicate increases in synthesis followed by breakdown without release. Regardless of these potential shortcomings, we have shown definitive evidence of hormone-dependent changes in the metabolic rate of monoamines in auditory areas.

## 5 | SUMMARY

In the present study, we tested predictions of a popular model of hormone-dependent salience of courtship signals.<sup>2,23,27,54,56,77</sup> According to the model, reproductive hormones act on monoaminergic modulatory inputs to auditory areas, which themselves respond to external events and encode behavioural relevance. In this way, auditory responses to courtship signals are amplified during the breeding season when they are most behaviourally relevant. Our results show partial support for this model. We did not find evidence of event-related monoaminergic responses that encode behavioural relevance. We did, however, find striking evidence that endocrine state modulates both event-related and event-independent monoaminergic activity in auditory areas. Because the salience and attractiveness of courtship signals is not determined solely by the auditory system<sup>27,54</sup>, future investigations should focus on how oestradiol shapes monoaminergic responses in other areas, such as the reward pathway and social decision-making network.<sup>2,56</sup>

## ACKNOWLEDGEMENTS

We thank Uyen Hoang, Brent Horton, Camden MacDowell and Michelle Park for technical assistance, as well as the Emory University Department of Biology for access to resources. This research was supported by an HHMI International Student Fellowship to CARS, NIH 5R01NS055125-04 to KWS, and NSF-IO5 0723805 to DLM.

## ORCID

D. L. Maney  <http://orcid.org/0000-0002-1006-2358>

C. A. Rodríguez-Saltos  <http://orcid.org/0000-0002-3162-7645>

## REFERENCES

- Eriksson D, Wallin L. Male bird song attracts females - a field experiment. *Behav Ecol Sociobiol.* 1986;19:297-299.
- Maney DL. The incentive salience of courtship vocalizations: hormone-mediated 'wanting' in the auditory system. *Hear Res.* 2013;305:19-30.
- Riebel K. Early exposure leads to repeatable preferences for male song in female zebra finches. *Proc R Soc Lond B Biol Sci.* 2000;267:2553-2558.
- Riebel K, Slater PJB. Testing female chaffinch song preferences by operant conditioning. *Anim Behav.* 1998;56:1443-1453.
- Maney DL, Cho E, Goode CT. Estrogen-dependent selectivity of genomic responses to birdsong. *Eur J Neurosci* 2006;23:1523-1529.
- Maney DL, Goode CT, Lange HS, Sanford SE, Solomon BL. Estradiol modulates neural responses to song in a seasonal songbird. *J Comp Neurol.* 2008;511:173-186.
- Moore M. Effect of female displays on endocrine physiology and behavior of male white-crowned sparrows, *Zonotrichia leucophrys.* *J Zool.* 1983;199:137-148.
- Henry KS, Lucas JR. Vocally correlated seasonal auditory variation in the house sparrow (*Passer domesticus*). *J Exp Biol.* 2009;212:3817-3822.
- Vélez A, Gall MD, Lucas JR. Seasonal plasticity in auditory processing of the envelope and temporal fine structure of sounds in three songbirds. *Anim Behav* 2015;103:53-63.
- Caras ML, Brenowitz E, Rubel EW. Peripheral auditory processing changes seasonally in Gambel's white-crowned sparrow. *J Comp Physiol A.* 2010;196:581-599.
- Yoder KM, Lu K, Vicario DS. Blocking estradiol synthesis affects memory for songs in auditory forebrain of male zebra finches. *NeuroReport.* 2012;23:922-926.
- De Groof G, Balthazart J, Cornil CA, Van der Linden. Topography and lateralized effect of acute aromatase inhibition on auditory processing in a seasonal songbird. *J Neurosci* 2017;37:4243-4254.
- De Groof G, Poirier C, George I, Hausberger M, Van der Linden A. Functional changes between seasons in the male songbird auditory forebrain. *Front Behav Neurosci.* 2013;7:1-13.
- Lattin CR, Stabile FA, Carson RE. Estradiol modulates neural response to conspecific and heterospecific song in female house sparrows: an in vivo positron emission tomography study. *PLoS ONE.* 2017;12:e0182875.
- Phillimore LS, Veysey AS, Roach SP. Zenk expression in auditory regions changes with breeding condition in male black-capped chickadees (*Poecile atricapillus*). *Behav Brain Res.* 2011;225:464-472.

16. Smeets WJA, Reiner A (eds). *Phylogeny and Development of Catecholamine Systems in the CNS of Vertebrates*. Cambridge: Cambridge University Press; 1994.
17. Schofield BR, Hurley L. Circuits for modulation of auditory function. In Oliver DL et al. eds. *The Mammalian Auditory Pathways, Springer Handbook of Auditory Research*. 2018;65:235-267.
18. Kitt CA, Brauth SE. Telencephalic projections from midbrain and isthmal cell groups in the pigeon. I. Locus coeruleus and subcoeruleus. *J Comp Neurol*. 1986;247:69-91.
19. Matragrano LL, Sanford SE, Salvante KG, Sockman KW, Maney DL. Estradiol-dependent catecholaminergic innervation of auditory areas in a seasonally breeding songbird. *Eur J Neurosci*. 2011;34:416-425.
20. Matragrano LL, Sanford SE, Salvante KG, Beaulieu M, Sockman KW, Maney DL. Estradiol-dependent modulation of serotonergic markers in auditory areas of a seasonally breeding songbird. *Behav Neurosci*. 2012;126:110-122.
21. Appeltants D, Del Negro C, Balthazart J. Noradrenergic control of auditory information processing in female canaries. *Behav Brain Res*. 2002;113:221-235.
22. Ritters LV, Pawlisch BA. Evidence that norepinephrine influences responses to male courtship song and activity within song control regions and the ventromedial nucleus of the hypothalamus in female European starlings. *Brain Res* 2007;1149:127-140.
23. Vyas A, Harding C, McGowan J, Snare R, Bogdan D. Noradrenergic neurotoxin, N-(2-chloroethyl)-N-ethyl-2-bromobenzylamine hydrochloride (DSP-4), treatment eliminates estrogenic effects on song responsiveness in female zebra finches (*Taeniopygia guttata*). *Behav Neurosci*. 2008;122:1148-1157.
24. Pawlisch BA, Stevenson SA, Ritters LV.  $\alpha$ 1-Noradrenergic receptor antagonism disrupts female songbird responses to male song. *Neurosci Lett*. 2011;496:20-24.
25. Pawlisch BA, Ritters LV. Selective behavioral responses to male song are affected by the dopamine agonist GBR-12909 in female European starlings (*Sturnus vulgaris*). *Brain Res*. 2010;1353:113-124.
26. Dai JB, Chen Y, Sakata JT. EGR-1 expression in catecholamine-synthesizing neurons reflects auditory learning and correlates with responses in auditory processing areas. *Neuroscience*. 2018;379:415-427.
27. Lynch KS. Understanding female receiver psychology in reproductive contexts. *Integr Comp Biol*. 2017;57:797-807.
28. LeBlanc MM, Goode CT, MacDougall-Shackleton EA, Maney DL. Estradiol modulates brainstem catecholaminergic cell groups and projections to the auditory forebrain in a female songbird. *Brain Res*. 2007;1171:93-103.
29. Grace AA, Floresco SB, Goto Y, Lodge DJ. Regulation of firing of dopaminergic neurons and control of goal-directed behaviors. *Trends Neurosci*. 2007;30:220-227.
30. Matragrano LL, Beaulieu M, Phillip JO, et al. Rapid effects of hearing song on catecholaminergic activity in the songbird auditory pathway. *PLoS ONE*. 2012;7:e39388.
31. Wolfson A. Regulation of refractory period in the photoperiodic responses of the white-throated sparrow. *J Exp Zool*. 1958;139:349-379.
32. Shank MC. The natural termination of the refractory period in the slate-colored junco and in the white-throated sparrow. *Auk*. 1959;76:44-54.
33. Griffiths R, Double MC, Orr K, Dawson RJG. A DNA test to sex most birds. *Mol Ecol*. 1998;7:1071-1075.
34. Maney DL, Lange HS, Raees MQ, Reid AE, Sanford SE. Behavioral phenotypes persist after gonadal steroid manipulation in white-throated sparrows. *Horm Behav*. 2009;55:113-120.
35. Pyle P, Howell SN, DeSante D, Yunick R. *Identification Guide to North American Birds*. Bolinas, CA: Slate Creek Press; 1997.
36. Thomas JW, Cáceres M, Lowman JJ, et al. The chromosomal polymorphism linked to variation in social behavior in the white-throated sparrow (*Zonotrichia albicollis*) is a complex rearrangement and suppressor of recombination. *Genetics*. 2008;179:1455-1468.
37. Lake JI, Lange HS, O'Brien S, Sanford SE, Maney DL. Activity of the hypothalamic-pituitary-gonadal axis differs between behavioral phenotypes in female white-throated sparrows (*Zonotrichia albicollis*). *Gen Comp Endocrinol*. 2008;156:426-433.
38. Horton BM, Moore IT, Maney DL. New insights into the hormonal and behavioural correlates of polymorphism in white-throated sparrows. *Anim Behav*. 2014;93:207-219.
39. Maney DL, Pinaud R. Estradiol-dependent modulation of auditory processing and selectivity in songbirds. *Front Neuroendocrinol*. 2011;32:287-302.
40. Sanford SE, Lange HS, Maney DL. Topography of estradiol-modulated genomic responses in the songbird auditory forebrain. *Dev Neurobiol*. 2010;70:73-86.
41. Matragrano LL, LeBlanc MM, Chitrapu A, Blanton ZE, Maney DL. Testosterone modulates genomic responses to song and monoaminergic innervation of auditory areas in a seasonally breeding songbird. *Dev Neurobiol*. 2013;73:455-468.
42. Sueur J, Aubin T, Simonis C. Seewave: a free modular tool for sound analysis and synthesis. *Bioacoustics*. 2008;18:213-226.
43. Ligges U, Krey S, Mersmann O, Schnackenberg S. tuneR: Analysis of music. 2013; R package version 1.2. 1.
44. Palkovits M, Brownstein MJ. Microdissection of brain areas by the punch technique. In: Cuello AC, ed. *Brain Microdissection Techniques*. New York, NY: Wiley;1983:2:1-36.
45. Vates GE, Broome BM, Mello CV, Nottebohm F. Auditory pathways of caudal telencephalon and their relation to the song system of adult male zebra finches (*Taeniopygia guttata*). *J Comp Neurol*. 1996;366:613-642.
46. Calabrese A, Woolley SM. Coding principles of the canonical cortical microcircuit in the avian brain. *Proc Natl Acad Sci USA*. 2015;112:3517-3522.
47. Kiltz CD, Breese GR, Mailman RB. Simultaneous quantification of dopamine, 5-hydroxytryptamine and four metabolically related compounds by means of reversed-phase high-performance liquid chromatography with electrochemical detection. *J Chromatogr B*. 1981;225:347-357.
48. Sockman KW, Salvante KG. The integration of song environment by catecholaminergic systems innervating the auditory telencephalon of adult female European starlings. *Dev Neurobiol*. 2008;68:656-668.
49. Barclay SR, Harding CF. Androstenedione modulation of monoamine levels and turnover in hypothalamic and vocal control nuclei in the male zebra finch: steroid effects on brain monoamines. *Brain Res*. 1988;459:333-343.
50. Barclay SR, Harding CF. Differential modulation of monoamine levels and turnover rates by estrogen and/or androgen in hypothalamic and vocal control nuclei of male zebra finches. *Brain Res*. 1990;523:251-262.
51. Harding CF, Barclay SR, Waterman SA. Changes in catecholamine levels and turnover rates in hypothalamic, vocal control, and auditory nuclei in male zebra finches during development. *J Neurobiol*. 1998;34:329-346.
52. Schielzeth H, Forstmeier W. Conclusions beyond support: overconfident estimates in mixed models. *Behav Ecol*. 2009;20:416-420.
53. Forlano PM, Licorish RR, Ghahramani ZN, et al. Attention and simulated motivated response to male advertisement call activates forebrain dopaminergic and social decision-making network nuclei in female midshipman fish. *Integr Comp Biol* 2017;57:820-834.
54. Maney DL, Rodriguez-Saltos CA. Hormones and the incentive salience of bird song; In: Bass AH, Sisneros JA, Fay RR, Popper AN,

- eds. *Hearing and Hormones. Springer Handbook of Auditory Research*. 2016;57:101-132.
55. Sockman KW, Lyons SM. How song experience affects female mate-choice, male song, and monoaminergic activity in the auditory telencephalon in Lincoln's sparrows. *Integr Comp Biol* 2017;57:891-901.
  56. Spool JA, Ritters LV. Associations between environmental resources and the "wanting" and "liking" of male song in female songbirds. *Integr Comp Biol*. 2017;57:835-845.
  57. Kruse AA, Stripling R, Clayton DF. Context-specific habituation of the zenk gene response to song in adult zebra finches. *Neurobiol Learn Mem* 2004;82:99-108.
  58. McKenzie TL, Hernandez AM, MacDougall-Shackleton SA. Experience with songs in adulthood reduces song-induced gene expression in songbird auditory forebrain. *Neurobiol Learn Mem*. 2006;86:330-335.
  59. Mello C, Nottebohm F, Clayton D. Repeated exposure to one song leads to a rapid and persistent decline in an immediate early gene's response to that song in zebra finch telencephalon. *J Neurosci*. 1995;15:6919-6925.
  60. Maney DL, Goode CT, Ball GF. Transduction of a non-photic cue: from the auditory system to a neuroendocrine response? *J Ornithol*. 2007;148:527-538.
  61. Maney DL, MacDougall-Shackleton EA, MacDougall-Shackleton SA, Ball GF, Hahn TP. Immediate early gene response to hearing song correlates with receptive behavior and depends on dialect in a female songbird. *J Comp Physiol A*. 2003;189:667-674.
  62. Velho TA, Lu K, Ribeiro S, Pinaud R, Vicario D, Mello CV. Noradrenergic control of gene expression and long-term neuronal adaptation evoked by learned vocalizations in songbirds. *PLoS ONE*. 2012;7:e36276.
  63. Lynch KS, Ball GF. Noradrenergic deficits alter processing of communication signals in female songbirds. *Brain Behav Evol* 2008;72:207-214.
  64. Lynch KS, Diekamp B, Ball GF. Colocalization of immediate early genes in catecholamine cells after song exposure in female zebra finches (*Taeniopygia guttata*). *Brain Behav Evol* 2012;79:252-260.
  65. Chen Y, Matheson LE, Sakata JT. Mechanisms underlying the social enhancement of vocal learning in songbirds. *Proc Natl Acad Sci USA*. 2016;113:6641-6646.
  66. Nordeen EJ, Holtzman DA, Nordeen KW. Increased Fos expression among midbrain dopaminergic cell groups during birdsong tutoring. *Eur J Neurosci*. 2009;30:662-670.
  67. Bharati IS, Goodson JL. Fos responses of dopamine neurons to socio-sexual stimuli in male zebra finches. *Neuroscience*. 2006;143:661-670.
  68. Clayton DF. The genomic action potential. *Neurobiol Learn Mem*. 2000;74:185-216.
  69. Gale SD, Perkel DJ. A basal ganglia pathway drives selective auditory responses in songbird dopaminergic neurons via disinhibition. *J Neurosci*. 2010;30:1027-1037.
  70. Ritters LV. Pleasure seeking and birdsong. *Neurosci Biobehav Rev*. 2011;35:1837-1845.
  71. Ritters LV, Olesen KM, Auger CJ. Evidence that female endocrine state influences catecholamine responses to male courtship song in European starlings. *Gen Comp Endocrinol*. 2007;154:137-149.
  72. Grace AA. Phasic versus tonic dopamine release and the modulation of dopamine system responsivity: a hypothesis for the etiology of schizophrenia. *Neuroscience*. 1991;41:1-24.
  73. Ikeda MZ, Jeon SD, Cowell RA, Remage-Healey L. Norepinephrine modulates coding of complex vocalizations in the songbird auditory cortex independent of local neuroestrogen synthesis. *J Neurosci*. 2015;35:9356-9368.
  74. Smith AR, Garris PA, Casto JM. Real-time monitoring of electrically evoked catecholamine signals in the songbird striatum using in vivo fast-scan cyclic voltammetry. *J Chem Neuroanat*. 2015;66:28-39.
  75. Hanson JL, Hurley LM. Context-dependent fluctuation of serotonin in the auditory midbrain. the influence of sex, reproductive state and experience. *J Exp Biol*. 2014;217:526-535.
  76. Keesom SM, Hurley LM. Socially induced serotonergic fluctuations in the male auditory midbrain correlate with female behavior during courtship. *J Neurophysiol*. 2016;115:1786-1796.
  77. Petersen CL, Hurley LM. Putting it in context: linking auditory processing with social behavior circuits in the vertebrate brain. *Integr Comp Biol*. 2017;57:865-877.

## SUPPORTING INFORMATION

Additional supporting information may be found online in the Supporting Information section at the end of the article.

**How to cite this article:** Rodríguez-Saltos CA, Lyons SM, Sockman KW, Maney DL. Sound-induced monoaminergic turnover in the auditory forebrain depends on endocrine state in a seasonally-breeding songbird. *J Neuroendocrinol*. 2018;30:e12606. <https://doi.org/10.1111/jne.12606>